


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 401/06, A61K 31/415	AI	(11) International Publication Number: WO 93/12107 (43) International Publication Date: 24 June 1993 (24.06.93)
(21) International Application Number: PCT/US92/10698 (22) International Filing Date: 16 December 1992 (16.12.92) (30) Priority data: 07/810,651 18 December 1991 (18.12.91) US (60) Parent Application or Grant (63) Related by Continuation US 07/810,651 (CIP) Filed on 18 December 1991 (18.12.91) (71) Applicant (for all designated States except US): SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033 (US).	(72) Inventors; and (75) Inventors/Applicants (for US only) : SHIH, Neng-Yang [US/US]; One Maple Drive, North Caldwell, NJ 07006 (US). GREEN, Michael, J. [US/US]; 43 Meadow Run Drive, Skillman, NJ 08558 (US). (74) Agents: JEANETTE, Henry, C. et al.; Schering-Plough Corporation, Patent Department, M3W, One Giralda Farms, Madison, NJ 07940-1000 (US). (81) Designated States: AU, BB, BG, BR, CA, CS, FI, HU, JP, KR, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG). Published With international search report.	

(54) Title: IMIDAZOYLALKYL SUBSTITUTED WITH A SIX MEMBERED NITROGEN CONTAINING HETEROCYCLIC RING



(57) Abstract

Disclosed is a compound of formula (1.0) or a pharmaceutically acceptable salt or solvate thereof. Also disclosed are pharmaceutical compositions comprising a pharmaceutically acceptable carrier and an effective amount of a compound of formula (1.0). Further disclosed is a method of treating allergy (for example asthma), inflammation, hypertension, raised intraocular pressure (such as glaucoma), i.e., a method of lowering intraocular pressure, sleeping disorders, states of hyper and hypo motility and acidic secretion of the gastrointestinal tract, hypo and hyperactivity of the central nervous system (for example, agitation and depression) and other CNS disorders (such as Alzheimers, Schizophrenia, and migraine) comprising administering an effective amount of a compound of formula (1.0) to a patient in need of such treatment.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CC	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MD	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

5


**IMIDAZOYLALKYL SUBSTITUTED WITH A SIX MEMBERED
NITROGEN CONTAINING HETEROCYCLIC RING**

10

BACKGROUND

H₃ receptor sites are known and are of current interest to
those skilled in the art—for example, see: West, Jr. et al., "Biexponential
Kinetics of (R)- α -[³H]Methylhistamine Binding to the Rat Brain H₃
Histamine Receptor", Journal of Neurochemistry, Vol. 55, No. 5, pp.
1612-1616, 1990; West, Jr. et al., "Identification of Two H₃-Histamine
Receptor Subtypes", Molecular Pharmacology, 38:610-613; and Korte et
al., "Characterization and Tissue Distribution of H₃ Histamine Receptors
in Guinea Pigs by N α -Methylhistamine", Biochemical and Biophysical
Research Communications, Vol. 168, No. 3, pp. 979-986..

Arrang et al. in U.S. 4,767, 778 (Issued August 30, 1988)
disclose a pharmaceutical composition containing a histamine
derivative of the formula:



wherein each of R₁, R₂, and R₄, represents a hydrogen or a methyl, or
R₁ and R₂ taken together represent a methylene, and R₃ is a hydrogen,
a methyl or a carboxy, with the proviso that R₁, R₂, R₃, and R₄ are not
simultaneously methyl groups. It is disclosed that the derivatives


30

SUBSTITUTE SHEET


- 2 -

behave as complete agonists of the H₃ receptors in rat brain and produce a maximal inhibition of release identical to that induced by histamine (approximately 60%). It is also disclosed that the histamine derivatives powerfully inhibit the release and synthesis of histamine by very selectively stimulating the H₃ receptors. Consequently, according to Arrang et al., the derivatives are likely to decrease histaminergic transmission in the digestive tract and in the nervous, cardiovascular and immune systems. Arrang et al. disclose that the derivatives can be used in therapy as a drug having sedative effects, as a sleep regulator, anticonvulsant, regulator of hypothalamo-hypophyseal secretion, antidepressant, and modulator of cerebral circulation. According to Arrang et al., inhibition of the release of inflammation messengers in various allergic conditions (e.g., asthma) is expected to result from stimulation of the H₃ receptors of the lung. It is further disclosed that the inhibition of release of gastric histamine is likely to exert antisecretory and antiulcerative effects. According to Arrang et al., modification of release of the messengers of immune responses is likely to modulate the latter responses.

EP 0 338 939 discloses compounds of the formula:




Derwent abstract 86-273706/42 for EP 0 197 840 discloses imidazole derivatives of the formula:




- 3 -

wherein R_1 is H, methyl or ethyl; R is H or R_2 ; and R_2 is 1-6C alkyl, piperonyl, 3-(benzimidazol-1-yl)propyl, -CZ-NHR₅ or a group (i):




5

wherein n is 0-3; X is a bond, O, S, NH, CO, CH=CH or a group (ii):



- 10 R_3 is H, methyl, halo, CN, CF₃ or COR₄; R_4 is 1-6C alkyl, 3-6C cycloalkyl or phenyl (optionally substituted by methyl or F); Z is O, S, NH, N-methyl or N-CN; and R_5 is 1-8C alkyl, 3-6C cycloalkyl (optionally substituted by phenyl), 3-6C cycloalkyl(1-3C)alkyl, phenyl (optionally substituted by methyl, halo or CF₃), phenyl(1-3C)alkyl; naphthyl, adamantyl or
- 15 p-toluenesulphonyl. It is disclosed that these compounds are psychotropic agents. It is also disclosed that these compounds antagonise the histamine H₃ receptors and increase the speed of cerebral histamine renewal.

- 20 Derwent abstract 90-184730/24 for U.S. 4,925,851 discloses 2- or 4-(2-(1H-imidazol-1-yl)ethyl) piperidine compounds useful as antitumour agents for inhibiting lymphoma, sarcoma, myeloma and leukaemia. The compounds have the formula:




25

wherein R is -CH₂(CH₂)_m-Me, -CO-(CH₂)_m-Me or -CO-CMe₂-R₂; m is 2-18; R_2 is H or Me; R_1 is -(CH₂)_n-R₃; n is 0-13; R_3 is H, i-Pr or t-Bu; and

- 4 -


the floating group is at the 2- or 4- position; with the proviso that (1) the sum of C atoms in R_1 does not exceed 13; and (2) the sum of C atoms in R and R_1 does not exceed 25.

Derwent abstract 90-180087/24 for EP 372125A discloses
5 compounds of the formula:



wherein X is O or S; R_1 is halo, CF_3 , CN, NO_2 , OH, or 1-6C alkoxy; R_2 is
10 H, 1-6C alkyl, aryl, 7-13C aralkyl, optionally substituted amino or 5- or
6-membered N-containing ring; and R_3 is 1-6C hydrocarbyl, 7-13C
aralkyl or 1-13C acyl. It is disclosed that these compounds have α_2 -
antagonist activity with no dopamine activity and that they are useful for
15 treating depression and other related illnesses (e.g., anxiety or cognitive
disorders).

Derwent abstract 88-309195/44 for U.S. 4935417
discloses compounds of the formula:




20


- 5 -

wherein (according to U.S. 4935417) R¹ is aryl, lower alkyl, cycloalkyl or hydrogen; R² is aryl, lower alkyl or hydrogen; R³ is lower alkyl, hydroxy or hydrogen; R⁴ is aryl or hydrogen; R⁵ is aryl or hydrogen; m is two or three; n is zero, one or two, provided that when R³ is hydroxy, n is one or two; and q is zero, one, two or three. U.S. 4935417 discloses that these compounds are calcium channel antagonists useful for treating mammals having a variety of disease states, such as stroke, epilepsy, hypertension, angina, migraine, arrhythmia, thrombosis, embolism and also for treatment of spinal injuries

10 Compounds known in the art include:



- 6 -




RN 81345-38-2
CA96(17):139642m

and

Known compounds in the art also include compounds of
the formula:

5



wherein R (Table 1) is:


- 7 -

TABLE 1


NO.	B	RN	CA
1	-CH ₃	106243-44-1	106(11):84602r
2	-CH(CH ₃) ₂	106243-45-2	106(11):84602r
3	H	106243-23-6	106(11):84602r
4	-C(S)NHC(CH ₃) ₂ CH ₂ C(CH ₃)	106243-93-0	106(11):84602r
5	-C(O)NHCH(CH ₃)(phenyl)	106243-90-7	---
6	-C(S)NH(p-chlorophenyl)	106243-85-0	---
7	-C(O)NH(phenyl)	106243-77-0	---
8	-C(NH)N(CH ₃)(cyclopropyl)	106243-73-6	---
9	-C(S)NHCH ₃	106243-61-2	---
10	-CH ₂ CH ₂ -phenyl	106243-49-6	---
11	-CH ₂ CH ₂ -p-fluorophenyl	106243-67-8	---
12	benzyl	106243-25-8	---

Additionally known compounds include:

5







RN 51746-84-0
CA80(15):82801a






RN 67319-35-1
CA89(13):109229v

- 8 -



5

- 9 -



5

In view of the art's interest in compounds which effect the H₃ receptors, novel compounds having agonist or antagonist activity on H₃ receptors would be a welcome contribution to the art. This invention provides just such a contribution by providing novel compounds having H₃ agonist or antagonist activity.


10

SUMMARY OF THE INVENTION

15

This invention provides compounds of the formula:

- 10 -



or a pharmaceutically acceptable salt or solvate thereof, wherein:

- 5 (A) m is an integer selected from the group consisting of: 1 and 2;
- (B) n and p are integers and are each independently selected from the group consisting of: 0, 1, 2, 3, and 4 such that the sum of n and p is 4 and T is a 6-membered ring;
- 10 (C) R³ and R⁴ are each independently bound to the same or different carbon atom of ring T such that there is only one R³ group and one R⁴ group in ring T, and each R¹, R², R³, and R⁴ is independently selected from the group consisting of:
- 15 (1) H;
- (2) C₁ to C₆ alkyl; and
- (3) -(CH₂)_q-R⁶ wherein q is an integer of: 1 to 7, and R⁶ is selected from the group consisting of: phenyl, substituted phenyl, -OR⁷, -C(O)OR⁷, -C(O)R⁷, -OC(O)R⁷, -C(O)NR⁷R⁸, CN and -SR⁷ wherein R⁷ and R⁸ are as defined below, and wherein the substituents on said substituted phenyl are each independently selected from the group consisting of:
- 20 -OH, -O-(C₁ to C₆)alkyl, halogen, C₁ to C₆ alkyl, -CF₃, -CN, and -NO₂, and wherein said substituted phenyl contains from 1 to 3 substituents;
- 25 (D) R⁵ is selected from the group consisting of:
- (1) H;
- (2) C₁ to C₂₀ alkyl;
- 30 (3) C₃ to C₆ cycloalkyl;

- 11 -

- 5 (4) $-\text{C}(\text{O})\text{OR}^{7'}$; wherein $\text{R}^{7'}$ is the same as R^7 defined below except that $\text{R}^{7'}$ is not H;
- (5) $-\text{C}(\text{O})\text{R}^7$;
- (6) $-\text{C}(\text{O})\text{NR}^7\text{R}^8$;
- (7) allyl;
- (8) propargyl; and
- (9) $-(\text{CH}_2)_q\text{R}^6$, wherein q and R^6 are as defined above, and when q is equal to 1, then R^6 is not OH or SH;
- 10 (E) R^7 and R^8 are each independently selected from the group consisting of: H, C_1 to C_6 alkyl, and C_3 to C_6 cycloalkyl;
- (F) the dotted line (-----) represents a double bond that is optionally present when m is 1, and n is not 0, and p is not 0 (i.e., the nitrogen in the ring is not bound directly to the carbon atom bearing the double bond), and when said
- 15 double bond is present then R^2 is absent; and
- (G) when m is 2, each R^1 is the same or different substituent for each m , and each R^2 is the same or different substituent for each m , and at least two of the substituents R^1 and/or R^2 are H.

20

Those skilled in the art will appreciate that the total number of substituents on each of the $-(\text{C})_n-$ and $-(\text{C})_p-$ groups is two, and that such substituents are independently selected from the group consisting of hydrogen, R^3 and R^4 , such that there is a total of only one R^3 and one

25 R^4 substituent in ring T.

This invention also provides pharmaceutical compositions comprising a pharmaceutically acceptable carrier and an effective amount of a Compound of Formula 1.0.

30 This invention further provides a method of treating allergy, (for example asthma), inflammation, hypertension, states of hyper and hypo motility and acidic secretion of the gastrointestinal tract, hypo and hyperactivity of the central nervous system (for example, agitation and depression) and other CNS disorders (such as Alzheimers, Schizophrenia, and migraine) comprising administering an effective

- 12 -

amount of a compound of Formula 1.0 to a patient in need of such treatment.

DETAILED DESCRIPTION OF THE INVENTION

5

As used herein the following terms have the following meanings unless indicated otherwise:

alkyl - represents a straight or branched, saturated hydrocarbon chain having from 1 to 20 carbon atoms;

10

cycloalkyl - represents a saturated carbocyclic ring having from 3 to 6 carbon atoms;

halogen (halo) - represents fluoro, chloro, bromo or iodo;

DMF - stands for (N, N-dimethylformamide);

LDA - stands for lithium diisopropylamide;

15

PDC - stands for pyridinium dichromate;

proton sponge - stands for [1,8-bis(dimethylamino)-naphthalene, N,N,N',N'-tetramethyl-1,8-naphthalenediamine];

SEM - stands for 2-(trimethylsilyl)ethoxymethyl;

THF - stands for tetrahydrofuran; and

20

TMEDA - stands for N,N,N',N'-tetramethylethylenediamine.



Also, unless stated otherwise, the substituents for the various embodiments described below are as defined for Formula 1.0.

Preferably, for compounds of Formula 1.0, m is 1; R⁵ is selected from the group consisting of H and C₁ to C₁₅ alkyl; and R¹ to R⁴ are each independently selected from the group consisting of: H, C₁ to C₆ alkyl, and -(CH₂)_q-R⁶ wherein R⁶ is phenyl. Most preferably, R⁵ is selected from the group consisting of H and C₁ to C₆ alkyl with H and methyl being even more preferable; and R³ and R⁴ are each independently selected from the group consisting of: H and methyl.


30

Representative compounds of this invention include compounds of the formula:

- 13 -





; and





5





Representative compounds of Formula 1.0 include compounds of the Formula:







10




- 14 -




5



- 15 -




5 Representative compounds of Formula 1.0 also include compounds of the Formula:









10

- 16 -




wherein R^1 is selected from the group consisting of: H and C_1 to C_6 alkyl, with H or methyl being preferred, and R^5 is selected from the group consisting of: H, C_1 to C_6 alkyl, $-C(O)OR^7$, $-C(O)R^7$, $-C(O)NR^7R^8$, and $-(CH_2)_q-R^6$ wherein R^6 is phenyl, with H or methyl being preferred

Representative compounds of Formula 1.0 further include:



- 17 -



5 Certain compounds of the invention may exist in different isomeric (e.g., enantiomers and diastereoisomers) forms. The invention contemplates all such isomers both in pure form and in admixture, including racemic mixtures. Enol forms are also included.

10 The compounds of Formula 1.0 can exist in unsolvated as well as solvated forms, including hydrated forms, e.g., hemi-hydrate. In general, the solvated forms, with pharmaceutically acceptable solvents such as water, ethanol and the like are equivalent to the unsolvated forms for purposes of the invention.

15 Certain compounds of the invention will be acidic in nature, e.g. those compounds which possess a carboxyl or phenolic hydroxyl group. These compounds may form pharmaceutically acceptable salts. Examples of such salts may include sodium, potassium, calcium, aluminum, gold and silver salts. Also contemplated are salts formed with pharmaceutically acceptable amines such as ammonia, alkyl
20 amines, hydroxyalkylamines, N-methylglucamine and the like.

 Certain basic compounds of the invention also form pharmaceutically acceptable salts, e.g., acid addition salts. For example, the nitrogen atoms may form salts with acids. Examples of suitable acids for salt formation are hydrochloric, sulfuric, phosphoric,
25 acetic, citric, oxalic, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those in the art. The salts are prepared by contacting the free

- 18 -

base form with a sufficient amount of the desired acid to produce a salt in the conventional manner. The free base forms may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous sodium hydroxide, potassium carbonate, ammonia and sodium bicarbonate. The free base forms differ from their respective salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the acid and base salts are otherwise equivalent to their respective free base forms for purposes of the invention.

All such acid and base salts are intended to be pharmaceutically acceptable salts within the scope of the invention and all acid and base salts are considered equivalent to the free forms of the corresponding compounds for purposes of the invention.

The following processes may be employed to produce compounds of Formula 1.0. Unless stated otherwise, reactions are conducted at an appropriate temperature which allows the reaction to proceed at a reasonable rate to completion.


A. PREPARATION OF COMPOUNDS WHEREIN m IS 1, n IS 0 AND

20

p is 4

SCHEME 1

- 19 -




- In Step 1 of Scheme 1, piperidine is reacted with di-tert-butyl dicarbonate ($(\text{tBOC})_2\text{O}$) in an organic solvent optionally in the presence of an organic base. The reaction is conducted at a temperature of about 0 to about 30°C. Preferably, methylene chloride is used as the organic solvent, but other suitable organic solvents include DMF and the like. Triethylamine is used as the organic base. Other bases which can be used include 4-dimethylaminopyridine and the like.
- In compound (1), the t-BOC group is chosen as an activating group on nitrogen which increases the kinetic acidity of the α -proton such that a lithio salt would result upon treatment with a strong organic base (for example, Step 2). Other activating groups on nitrogen, known in the art

- 20 -

that can also be employed include nitroso, phosphoryl, hindered acyl, and formamidine. (See Aldrichimica Acta, Vol. 8, No. 3, 1985).

In Step 2 of Scheme 1, the anion of compound (1) is reacted with compound (2)


5



to produce compound (3). The reaction is conducted in an inert organic solvent containing an organic base and TMEDA (N,N,N',N'-tetramethylethylenediamine). The reaction is conducted at a temperature of about -78 to about 25°C (room temperature). Tetrahydrofuran is preferably used as the solvent, other suitable solvents include diethyl ether and the like. The anion of (1) is prepared by treatment of (1) with sec-butyllithium in THF at -78°C.

15

Z represents the protecting group




Z can be other protecting groups, such as 2-(trimethylsilyl)ethoxymethyl, benzyloxycarbonyl and the like; however, unless stated otherwise, Z preferably represents the trityl group in the process described below for making the compounds of this invention.

20

- 21 -

Those skilled in the art will appreciate that other protecting groups known in the art may be used—such as, for example, base sensitive groups wherein the protected compounds would be deprotected using basic conditions (e.g., NaOH). The processes described herein wherein the protected compound is deprotected under acidic conditions may also be carried out under basic conditions when a base sensitive protecting group is used.

Compound (2) is prepared in two steps from compound (4):



Compound (4) is reacted with an organometallic reagent R^1M , wherein M is Li or MgBr, to produce compound (5). The reaction takes place in an inert organic solvent at a temperature of about -78°C to 0°C . Suitable inert organic solvents include: THF, diethyl ether and the like. Compound (5) is then reacted with thionyl chloride in an inert organic solvent, such as benzene or CH_2Cl_2 , in the presence of base to generate compound (2). The reaction is conducted at a temperature of about -20°C to 80°C . Suitable bases include: pyridine, triethylamine, and the like. Preferably, triethylamine is used as the amine base. Compound (4) can be obtained by following the literature procedure set forth in J.K. Kelly, et al., J. Med. Chem., 20, 721 (1977).

In Step 3 of Scheme 1, compound (3) is treated with HCl or similar acid in an inert organic solvent such as ethyl acetate or dioxane, at a temperature of about 0°C to selectively deprotect (3) thus producing compound (6).

In Step 4 of Scheme 1, compound (6) may be reacted with (i) $\text{R}^5\text{-X}$ (when R^5 is $-\text{C}(\text{O})\text{R}^7$, $-\text{C}(\text{O})\text{OR}^7$, $-\text{C}(\text{O})\text{NR}^7\text{R}^8$ or alkyl) in an organic solvent optionally in the presence of a suitable base (e.g., triethylamine); or (ii) $\text{R}^5\text{A-CHO}$ (when R^5 is alkyl, cycloalkyl, allyl,

- 22 -

propargyl, benzyl or substituted benzyl) in the presence of $\text{NaBH}_3(\text{CN})$ (sodium cyanoborohydride) or other hydrogenating conditions (e.g. $\text{H}_2/\text{Pd}/\text{ROH}$) in an organic solvent; to produce compound (7). $\text{R}^{5\text{A}}$ represents an R^5 group that has one less $-\text{CH}_2-$ group. Preferably,
5 CH_2Cl_2 is used as the solvent when $\text{R}^5\text{-X}$ is used, and tetrahydrofuran is used as the solvent when $\text{R}^{5\text{A}}\text{-CHO}$ is used. X represents a suitable leaving group such as Cl, Br, I, or $-\text{OCH}_3$. The reaction ((i) or (ii)) can be performed at a temperature within the range of about -30 to about 80°C . Compound (7), when R^5 is $-\text{C}(\text{O})\text{NR}^7\text{H}$, may be prepared by
10 reacting compound (6) with $\text{O}=\text{C}=\text{N}-\text{R}^7$ in an organic solvent, such as CH_3CN or toluene. The reaction is performed at a temperature in the range of about 20 to reflux. Compound (6), or compound (7) wherein R^5 is $-\text{C}(\text{O})\text{O}(\text{t-butyl})$, can be reacted with aqueous acid (e.g., HCl, HBr, and the like), at a temperature of about 25 to about 100°C , to produce
15 compound (8) wherein R^5 is H.


Compound (7), wherein R^5 is $-\text{C}(\text{O})\text{OR}^7$, may be reacted with NHR^7R^8 in an organic solvent at a temperature of about 25 to about 100°C to form a compound of (7) wherein R^5 is $-\text{C}(\text{O})\text{NR}^7\text{R}^8$. The suitable organic solvents include THF, toluene, DMF and the like.

20 In Step 5 of Scheme 1, compound (7) may be deprotected by treatment with dilute aqueous acid, such as HCl or HBr, at a temperature of about 70 to about 90°C to produce compound (8). Other protecting groups are removed by methods well known in the art.

Alternatively compound (3) of Preparation A can be
25 prepared from compounds (1) and (4) according to Scheme 2:

- 23 -

SCHEME 2




5 In Step 1 of Scheme 2, compound (1)--see Step 1 of Scheme 1--is reacted with compound (4) in accordance with the procedure set forth in Step 2 of Scheme 1.

10 In Step 2 of Scheme 2, compound (9) is oxidized to produce compound (10). The oxidation is accomplished by treating compound (9) with an oxidizing agent, such as MnO_2 or PDC (pyridinium dichromate), in an organic solvent, such as tetrahydrofuran or methylene chloride, at a temperature of about 20 to about 80°C.

In Step 3 of Scheme 2, compound (10) is reacted, under usual Wittig reaction conditions, with compound (11)

- 24 -



5 in an organic solvent at a temperature of about 25 to about 80°C to produce compound (12). Preferably, the organic solvent is tetrahydrofuran; however, other suitable solvents, such as 1,4-dioxane and the like, can be used. In compound (11), R^{1A} represents an R¹ group which has one less -CH₂- group.


10 In Step 4 of Scheme 2, compound (3) is produced when compound (12) is hydrogenated in THF with H₂ using a Pd-C (palladium/carbon) catalyst. Other organic solvents which can be used include ethyl acetate, methanol and the like. Other suitable metals such as Pt, Pd-Al₂O₃, Ra-Ni, NiB, and Pd-CaCO₂ can also be employed as the hydrogenation catalyst.

15 In all the preparations that follow, intermediate compounds wherein the imidazole nitrogen is protected by Z and the nitrogen of the cyclic six membered amine is substituted with -C(O)O(t-butyl) or unsubstituted, i.e., hydrogen is bound to the amine nitrogen, such as in compounds (7) or (6), respectively, such intermediate compounds can
20 be reacted with aqueous acid (e.g., HCl, HBr, and the like), at a temperature of about 25 to about 100°C, to produce deprotected final products wherein R⁵ is H, e.g., compound (8).

25 B. PREPARATION OF COMPOUNDS WHEREIN m IS 1, n IS 1 AND p IS 3

SCHEME 3

- 25 -



- In Step 1 of Scheme 3, α -valerolactam is reacted with R^9-L to place the R^9 on the indicated nitrogen atom in compound (13). R^9 can either be $Si(CH_3)_2C(CH_3)_3$ or other protecting group, or R^9 can be an alkyl, cycloalkyl, benzyl, substituted benzyl, allyl, or propargyl group. In the latter, R^9 is the same as R^5 . L is a leaving group, such as Cl , Br , I or $-OSO_2CF_3$. The reaction is conducted in an

- 26 -

organic solvent such as THF, diethyl ether or 1,4-dioxane in the presence of a suitable base, such as lithium diisopropylamide (LDA), KH or NaH. The reaction takes place at a temperature within the range of about -78 to about 80°C.

5 In Step 2 of Scheme 3, the anion of compound (13) is reacted with compound (4) in an organic solvent, containing an organic base, at a temperature of about -78 to about 25°C to produce compound (14). Suitable organic solvents include tetrahydrofuran and the like. Preferably, the organic base used to generate the anion of (13) is lithium
10 diisopropylamide or $MN[Si(CH_3)_3]_2$ wherein M is a metal cation such as Li, Na, or K. Z is trityl or other suitable protecting group.

In Step 3 of Scheme 3, compound (14) is reacted with R^1-Q , wherein Q is Li or MgBr, in tetrahydrofuran containing CuCN and a Lewis acid, such as $BF_3 \cdot (C_2H_5)_2O$, $(CH_3)_3SiCl$ and the like, to
15 produce compound (15). The reaction is conducted at a temperature of about -78 to about 20°C. Tetrahydrofuran is the preferred organic solvent; however, other suitable solvents include diethyl ether and the like.

Compound (15) in Scheme 3 can alternatively be prepared
20 by reacting an anion of compound (13) with compound (2) (see Step 2 of Scheme 1) in the presence of an organic base. Inert organic solvents, such as THF, 1,4-dioxane and the like can be employed. Preferably, the organic base used to generate the anion of (13) is lithium diisopropylamide or $MN[Si(CH_3)_3]_2$ wherein M is a metal cation such as
25 Li, Na, or K. The reaction takes place at a temperature of about -78°C to about 25°C.

In Step 4 of Scheme 3, the enolate of compound (15) is reacted with R^4-L in an inert organic solvent to produce compound (16). The reaction can be conducted at a temperature of about 0 to about
30 50°C. L is a suitable leaving group such as Cl, Br, I and the like. Preferably, KH is used as the base to form the enolate, but other suitable bases include LDA and the like. Suitable organic solvents include THF, 1,4-dioxane and the like. Preferably, THF is used.

- 27 -

In Step 5 of Scheme 3, compound (16) is reduced with LiAlH_4 at a temperature of about 0 to about 65°C to produce compound (17). The reduction is conducted in tetrahydrofuran, 1,4-dioxane and the like; however, THF is preferable. Other suitable reducing agents include BH_3 (borane) and the like.

Step 6 of Scheme 3 (compound (17) to (19)), is followed when R^9 is the desired substituent on the nitrogen, such as alkyl, cycloalkyl, benzyl, substituted benzyl, allyl, or propargyl. Step 7 of Scheme 3 (compound (17) to (18)), is followed when R^9 is $-\text{Si}(\text{CH}_3)_2\text{C}(\text{CH}_3)_3$.


In Step 6 of Scheme 3, compound (17), when R^9 is alkyl, cycloalkyl, benzyl, substituted benzyl, allyl or propargyl, is deprotected following the procedure in Step 5 of Scheme 1 to produce compound (19). Alternatively, in Step 7 of Scheme 3, compound (17) is treated with tetrabutylammonium fluoride in tetrahydrofuran at a temperature of about 0 to about 50°C to produce compound (18).

In Step 8 of Scheme 3, the procedures in Steps 4 and 5 of Scheme 1 are followed so that compound (18) is converted to compound (19).

C. PREPARATION OF COMPOUNDS WHEREIN m IS 1, n IS 1 AND p IS 3

SCHEME 4

- 28 -




- Compound (20) is reacted with R³-M (wherein M is Li, ZnBr or MgBr) in tetrahydrofuran containing BF₃·(C₂H₅)₂O and CuCN at a temperature of about -78 to about 20°C to produce compound (21). Other suitable solvents such as diethyl ether can be used. Compound (21) is then converted to compound (22) in accordance with the reaction steps set forth in Scheme 3.

10 D. PREPARATION OF COMPOUNDS WHEREIN m IS 1, n IS 2 AND p IS 2

SCHEME 5

- 29 -



In Step 1 of Scheme 5, compound (23), wherein R is CH₃ or CH₂-C₆H₅, is reacted with Mg in an inert organic solvent, such as THF, at a temperature of about 50 to about 70°C to form the Grignard reagent. The Grignard reagent is then coupled with compound (2), in the presence of CuI or CuCN, to produce compound (24). This coupling reaction is conducted at a temperature of about -78 to about 20°C. THF is the preferable solvent; however, other solvents such as 1,4-dioxane and diethyl ether can be used.

In Step 2 of Scheme 5, compound (24), wherein R is CH₃, is demethylated to produce compound (25) by following the literature procedures set forth either in the J. Am. Chem. Soc., 110, 8256 (1988) or in the J. Org. Chem., 49, 2081 (1984). For example, the latter procedure compound (24) is reacted with ClC(O)OCHClCH₃ in 1,2-dichloroethane at about 0 to about 85°C in the presence of proton sponge to produce compound (25). Alternatively, compound (24), wherein R is CH₂-C₆H₅, is converted to compound (25) under either the hydrogenation conditions or transfer-hydrogenation conditions. For the former, the reaction is conducted in THF using Pd-C as catalyst, at a temperature of about 20 to about 70°C. Other organic solvents which can be used

- 30 -

include ethanol and the like. Other suitable metals such as Pt, Pd-Al₂O₃ and the like can also be employed as catalysts. For the latter, the source of hydrogen is ammonium formate (NH₄COO), the catalyst is Pd/C, at a temperature of about 20 to about 80°C. The suitable organic solvents include methanol, ethanol and the like.


In Step 3 of Scheme 5, compound (25) is converted to compound (26) in accordance with the procedure set forth in Step 4 and 5 of Scheme 1.

Compound (26), wherein R⁵ is CH₃ or CH₂-C₆H₅, can be directly prepared from compound (24), by a simple deprotection of the Z group, by following the procedure set forth in Step 5 of Scheme 1.

E. PREPARATION OF COMPOUNDS WHEREIN m IS 1, n IS 2 AND
p is 2

SCHEME 6

- 31 -



In Step 1 of Scheme 6, compound (2) is reacted with a metal ($M=Zn$ or Mg) in an inert organic solvent, such as THF and the like, at a temperature of about -20 to about $60^\circ C$, to generate either an organozinc reagent (when $M=Zn$) or a Grignard reagent (when $M=Mg$). This organometallic reagent is then reacted with $CuCN \cdot 2LiCl$ in THF to form an intermediate, compound (27), which is subsequently coupled with compound (20), in the presence of $BF_3 \cdot (C_2H_5O)_2$ or $(CH_3)_3SiCl$, to form compound (28). This coupling reaction is conducted in an inert organic solvent, such as THF, 1,4-dioxane or diethyl ether, at a temperature of about -78 to $50^\circ C$.

In Step 2 of Scheme 6, the anion of compound (28) is reacted with R^3-X and then with R^4-X to produce compound (29). X represents a suitable leaving group, such as Cl, Br, I or $-OSO_2-CF_3$.

- 32 -

Each reaction to place each substituent group on the ring takes place in an organic solvent using an organic base. THF is the solvent usually used; however, other suitable solvents include 1,4-dioxane, diethyl ether and the like. Examples of organic bases include lithium diisopropyl-
5 amide, $MN[Si(CH_3)_3]_2$, KH and like. M represents a suitable metal cation such as Na, Li, K, and the like. The reaction is usually conducted at a temperature of about -78 to about 80°C.


In Step 3, compound (29) is reduced to compound (30) in accordance with the procedure set forth in Step 5 of Scheme 3.

10 In Step 4, the procedure in Step 6 or Steps 7 and 8 of Scheme 3 can be followed so that compound (30) is converted to compound (31).

15 F. PREPARATION OF COMPOUNDS WHEREIN m IS 1, n IS 2 AND
p is 2

SCHEME 7

- 33 -



In Step 1 of Scheme 7, 4-chloropiperidine is reacted with R⁹-L to place an R⁹ on the nitrogen atom of compound (32). Both R⁹ and L are as defined above in Step 1 of Scheme 3. The reaction is conducted in an organic solvent, such as THF, methylene chloride and the like, in the presence of a suitable base, such as triethylamine or 4-dimethylamino pyridine. The reaction takes place at a temperature within the range of about 0 to about 70°C.

In Step 2 of Scheme 7, compound (32) is converted to compound (34) in accordance with the reaction conditions set forth for

- 34 -

the conversion of compound (23) to compound (24) (See Step 1 of Scheme 5).

5 In Step 3 of Scheme 7, compound (34) is chlorinated with sulfuryl chloride to form compound (35). This reaction is conducted at a temperature of about 0 to about 50°C in an inert organic solvent and in the presence of suitable base. Dichloromethane is the preferred solvent, other solvents such as 1,2-dichloroethane can also be used. Suitable organic bases include triethylamine, 4-dimethylaminopyridine and the like.


10 In Step 4 of Scheme 7, the conversion of compound (35) to compound (36) is achieved in a two step sequence. First, compound (35) is reacted with formamide at about 175°C (See Chem. Ber., Vol. 86, p. 88 (1953)) to form the imidazole ring. In the next step, the imidazole is protected with a Z group by reacting the product of the first step with Z-L
15 in the presence of an organic base to give compound (36). L is a leaving group, such as Cl or Br, and Z is as defined above in Step 2 of Scheme 1. Suitable organic solvents for the protection step include methylene chloride, dimethylformamide and the like. Suitable organic bases include triethylamine and the like. The reaction may be
20 performed at a temperature of about 0 to about 50°C.

In Step 5 of Scheme 7, compound (36) is converted to compound (26A) in accordance with the sequence set forth in Step 6 or Steps 7 and 8 in Scheme 3.

25 G. PREPARATION OF COMPOUNDS WHEREIN m IS 1, n IS 2 AND
p is 2

SCHEME 8

- 35 -



- 5 In Step 1 of Scheme 8, 4-halopyridine (wherein X is Cl, Br or I) is converted to the corresponding organocuprate reagent (37) by reaction with butyllithium and then with $\text{CuCN}(\text{CN})_2\text{LiCl}$ in an inert organic solvent at a temperature of about -78 to about 0°C . Suitable organic solvents include THF, diethylether and the like. The organocuprate (37) thus formed is then coupled with the α,β -unsaturated aldehyde (33) to form compound (38). The coupling reaction is performed at a temperature of

- 36 -

-78°C to 20°C in THF in accordance with the procedure set forth in Step 1 of Scheme 6.

In Step 2 of Scheme 8, compound (38) is reacted with sulfonyl chloride, in accordance with the procedure set forth in Step 3 of of Scheme 7, to give compound (39).


In Step 3 of Scheme 8, compound (39) is then converted to compound (40) using the Brederick Reaction conditions (See Chem. Ber., 86, 88 (1954)).

In Step 4 of Scheme 8, compound (40) is hydrogenated to compound (41), under a hydrogen pressure of about 1 to about 100 atmospheres, in the presence of suitable catalyst and in an acidic aqueous solvent. Suitable hydrogenation catalysts include Pd-black or Rh/C and the like. Suitable solvents include 10% aqueous hydrochloric acid, or 10% aqueous sulfuric acid, or the like. The reaction takes place at a temperature of about 20 to about 100°C.

In Step 5 of Scheme 8, compound (41) is converted to compound (26A) in accordance with the procedure set forth in Step 4 of Scheme 1.

H. PREPARATION OF COMPOUNDS WHEREIN m IS 1, n IS 1 AND
p is 3

SCHEME 9




By following the reaction sequence set forth in Scheme 8, compound (19A) is prepared by starting with the 3-halopyridine, wherein X is Cl, Br, or I.

I. PREPARATION OF COMPOUNDS WHEREIN m IS 1, n IS 0 AND
p is 4

SCHEME 10


5



2-Halopyridine, wherein X is Cl, Br or I, is converted to
10 compound (8A) by following the sequence set forth in Scheme 8.


J. PREPARATION OF COMPOUNDS WHEREIN m IS 2, n IS 0 AND p IS 4; OR m IS 2, n IS 1 AND p IS 3

15 By following the steps in Schemes 1 to 4 with the exception that compound (42)






20 is used instead of compound (2) and compound (43)

- 38 -



is used in place of compound (4), compounds (44), (45) and/or (47)




5

are produced. Compound (44) is produced by following Scheme 1 or Scheme 2, and compound (45) and (47) are produced by following Scheme 3 and Scheme 4, respectively. G represents a suitable leaving group such as Br, I, -OSO₂-C₆H₄-CH₃, -OSO₂-CF₃, -OSO₂-CH₃ and the like. The preparation of compound (42) is described below.

10

- 39 -

SCHEME 11 - PREPARATION OF COMPOUND (42)

- 5 Compound (48) is produced by reacting compound (46) (wherein Y is either CN or COOR¹⁰ wherein R¹⁰ is C₁ to C₅ alkyl) in an organic solvent containing an organic base with R¹-L and then with R²-L in accordance with the method set forth in Step 2 of Scheme 6. Preferably, the organic solvent is tetrahydrofuran and the organic base is lithium diisopropylamide. L is a suitable leaving group such as Cl, Br, I, -OSO₂-CF₃ and the like. Compound (48) is then reduced to compound (49) in THF at a temperature of about -78 to about 70°C using diisobutylaluminum hydride (when Y is CN) or bis(2-methoxyethoxy)aluminum hydride (when Y is COOCH₃; see for example
- 10 R. Kanazawa & T. Tokoroyama, *Synthesis*, 526 (1976)) as the reducing agent. Alternatively, the sequence of the preparation of compound (49) from (46) can be switched--i.e., reduction first and then introduction of R¹ and R².



- Compound (49) is then reacted with either lithium
- 20 aluminum hydride (when R¹ is H) or R¹-Q (when R¹ is not H) in tetrahydrofuran at a temperature of about -78 to about 0°C to produce compound (50). Q represents Li or MgBr. When G represents a halide

- 40 -

- (i.e., Cl, Br, or I), then compound (42) is produced by either reacting compound (50) with $(C_6H_5)_3P/CG_4$ or $(C_6H_5)_3PG_3$ (see Fiser & Fiser, Reagents for Organic Synthesis, Vol. 1, p1247 (1967)). When G represents $-OSO_2-C_6H_4-CH_3$, $-OSO_2-CH_3$ or $-OSO_2-CF_3$, then
- 5 compound (42) is produced by reacting compound (50) with $Cl-SO_2-C_6H_4-CH_3$, $Cl-SO_2-CH_3$ or $Cl-SO_2-CF_3$, respectively, in methylene chloride containing triethylamine (as base) at a temperature of about -78 to about $0^\circ C$.

10 K. PREPARATION OF COMPOUNDS WHEREIN m IS 2, n IS 2 AND p IS 2

- By following the steps in Scheme 5 and Scheme 6, with the exception that compound (42) (see Preparation J) is used instead of
- 15 compound (2), compound (51) is produced following Scheme 5, and compound (52) is produced following Scheme 6.




20 L. PREPARATION OF COMPOUNDS WHEREIN DOUBLE BOND INDICATED IN FORMULA 1.0 IS PRESENT AND WHEREIN m IS 1, n IS 2 AND p IS 2


SCHEME 12

25

- 41 -



In Step 1 of Scheme 12, compound (32) is reacted with Mg, in accordance with the procedure set forth in Step 1 of Scheme 5, to form the corresponding Grignard reagent. This Grignard reagent is reacted with compound (4)




to form compound (53). The reaction takes place in a suitable organic solvent at a temperature of about -78 to about 20°C. Suitable organic solvents include THF; 1,4-dioxane and the like. Preferably THF is used.

In Step 2 of Scheme 12, compound (53) is converted to compound (54), wherein R^{5'} is H, alkyl, cycloalkyl, benzyl, substituted benzyl, allyl or propargyl group, using an acidic aqueous solution, such as HCl, HBr or the like. The conversion takes place at a temperature of about 50 to about 100°C. Compound (54), wherein R^{5'} is H, is then converted to compound (55) in accordance with Step 4 of Scheme 1.

- 42 -

M. PREPARATION OF COMPOUNDS WHEREIN THE DOUBLE BOND IN FORMULA 1.0 IS PRESENT AND WHEREIN m IS 1, n IS 1 AND p IS 3

5

SCHEME 13

- 43 -


In Step 1 of Scheme 13, compound (57) is produced when compound (56) is reacted with (t-BOC)₂O and triethylamine. The reaction is conducted in an organic solvent, such as methylene chloride or DMF, using a temperature within the range of about 0 to about 25°C (room temperature).

In Step 2 of Scheme 13, compound (58) is produced by treating compound (57) with an oxidizing agent such as pyridinium dichromate or similar oxidizing agent. The oxidation reaction is conducted in an organic solvent, such as methylene chloride, using a temperature of about 25 to about 50°C.

In Step 3 of Scheme 13, compound (59) is produced when the enolate of compound (58) is reacted with R³-L wherein L is a suitable leaving group, such as halogen (e.g., Cl, Br, or I), -OSO₂CF₃ and the like. The reaction takes place in an organic solvent, such as tetrahydrofuran or benzene, containing a suitable base, such as NaH, KH, LDA, or LiN(Si(CH₃)₃)₂. Preferably, tetrahydrofuran is used as the solvent and LDA is used as the base. The reaction is conducted at a temperature of about 0 to about 80°C.

In Step 4 of Scheme 13, the anion of compound (59) is reacted with R⁴-L using the same procedure set forth in Step 3 of Scheme 13 in order to produce compound (60) and/or (61).


In Step 5 of Scheme 13, compound (63) or (63A) is obtained when compound (60) or (61) is reacted with compound (62)



The reaction takes place in an organic solvent, such as tetrahydrofuran, DMF or benzene, containing a suitable base, such as NaH, LDA, or LiN(Si(CH₃)₃)₂. Preferably, tetrahydrofuran is used as the solvent and LDA is used as the base. The reaction is conducted at a temperature of

- 44 -

about 0 to about 80°C. Compound (62) is obtained by reacting compound (2A)



5


with P(C₆H₅)₃ in an organic solvent, such as methylene chloride, CH₃CN, tetrahydrofuran and the like, using a temperature of about 25 to about 50°C. In compounds (62) and (2A), Z represents trityl or SEM.

10 In Step 6 of Scheme 13, compound (64) and (64A) are prepared from compound (63) and (63A), respectively, by following the same procedure set forth in Steps 3 to 5 of Scheme 1.

In the steps of Scheme 13, alkylations (i.e., Steps 3 and 4) are only if desired and R³ and R⁴ are as defined for Formula 1.0.

15 N. PREPARATION OF COMPOUNDS WHEREIN THE DOUBLE BOND IN FORMULA 1.0 IS PRESENT AND WHEREIN m IS 1, n IS 2 AND p IS 2



20 By following the steps in Scheme 13, with the exception that compound (65)



is used instead of compound (56), compound (66) and (66A)

25

- 45 -



is produced.

In the above processes, certain functional groups may be incompatible with some transformations described herein, and consequently it is sometimes desirable and/or necessary to protect certain groups during the reactions. Certain protecting groups are employed in the above processes but, as those skilled in the art will recognize, other protecting groups may be used in their place. Conventional protecting groups are operable as described in Greene, T.W., and Wuts, P.G.M., "Protective Groups In Organic Synthesis," John Wiley & Sons, New York, 1991; the disclosure of which is incorporated herein by reference thereto. After the reaction or reactions, the protecting groups may be removed by standard procedures.

The compounds of this invention are either agonists or antagonists of the histamine H₃ receptor. The binding affinity of the compounds of the invention to the H₃ receptor may be demonstrated by the procedure described below:

H₃ Receptor Binding Assay

The source of the H₃ receptors in this experiment was guinea pig brain. The animals used weighed 400-600 g. The tissue was homogenized using a Polytron in a solution of 50 mM Tris, pH 7.5. The final concentration of tissue in the homogenization buffer was 10% w/v. The homogenates were centrifuged at 1000 x g for 10 min. in order to remove clumps of tissue and debris. The resulting supernatants were then centrifuged at 50,000 x g for 20 min. in order to sediment the membranes, which were next washed 3 times in homogenization buffer

- 46 -






(50,000 x g for 20 min. each). The membranes were frozen and stored at -70°C until needed.

5 All compounds to be tested were dissolved in DMSO and then diluted into the binding buffer (50 mM Tris, pH 7.5) such that the final concentration was 2 µg/mL with 0.1% DMSO. Membranes were then added (400 µg of protein) to the reaction tubes. The reaction was started by the addition of 3 nM [³H]R-α-methylhistamine (8.8 Ci/mmol) or [³H]-N-methylhistamine (80 Ci/mmol) and incubated at 30° for 30 min. Bound ligand was separated from unbound ligand by filtration, and the
10 amount of radioactive ligand bound to the membranes was quantitated by liquid scintillation spectrometry. All incubations were performed in duplicate and the standard error was less than 10% in all instances. Compounds that inhibited greater than 70% of the specific binding of radioactive ligand to the receptor were serially diluted to determine a K_i
15 (µM). The results are given in Table 2.

In Table 2, the compound represented by (a*) is known in the art.

- 47 -

TABLE 2

COMPOUND	H ₃ Binding K _i (μ M)
 (a*)	0.014
 (6.2)	0.19
 (7.2)	0.0008
 (9.1)	0.012
 (7.3)	0.0003

- For preparing pharmaceutical compositions from the
- 5 compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. The powders and tablets may be comprised of from about 5 to about 70 percent active ingredient. Suitable solid carriers are

- 48 -

known in the art, e.g. magnesium carbonate, magnesium stearate, talc, sugar, lactose. Tablets, powders, cachets and capsules can be used as solid dosage forms suitable for oral administration.

For preparing suppositories, a low melting wax such as a
5 mixture of fatty acid glycerides or cocoa butter is first melted, and the active ingredient is dispersed homogeneously therein as by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool and thereby solidify.

Liquid form preparations include solutions, suspensions
10 and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection.

Liquid form preparations may also include solutions for intranasal administration.

Aerosol preparations suitable for inhalation may include
15 solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier, such as an inert compressed gas.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include
20 solutions, suspensions and emulsions.

The compounds of the invention may also be deliverable transdermally. The transdermal compositions can take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in
25 the art for this purpose.

Preferably the compound is administered orally.

Preferably, the pharmaceutical preparation is in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component, e.g., an
30 effective amount to achieve the desired purpose.

The quantity of active compound in a unit dose of preparation may be varied or adjusted from about 0.1 mg to 1000 mg, more preferably from about 1 mg to 500 mg, according to the particular application.


- 49 -

The actual dosage employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired.

The amount and frequency of administration of the compounds of the invention and the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated. A typical recommended dosage regimen is oral administration of from 1 mg to 2000 mg/day preferably 10 to 1000 mg/day, in one to four divided doses to achieve relief of the symptoms. The compounds are non-toxic when administered within this dosage range.

The invention disclosed herein is exemplified by the following preparative examples, which should not be construed to limit the scope of the disclosure. Alternative mechanistic pathways and analogous structures within the scope of the invention may be apparent to those skilled in the art.


25

EXAMPLE 1

To a solution of diisopropylamine (1.078 mL) in anhydrous THF (7 mL) was added n-butyllithium (3.08 mL; 2.5M) dropwise at 0°C.


- 50 -

The resulting solution was stirred at 0°C for 40 minutes and then was cooled to -23°C. To this mixture was added N-methyl-2-piperidinone (0.80 mL) (1), the mixture was stirred at -23°C for 0.5 hour, and then at -78°C for 1 hour. To the above mixture was added dropwise a solution of 4-chloromethyl-N-trityl-imidazole (2.70 g)



in anhydrous THF (14 mL). The mixture was stirred at -78°C for 4 hours and then was allowed to warm up to room temperature slowly overnight (16 hours). Water and ethyl acetate were added to the mixture, the resulting mixture was shaken vigorously, the layers separated, and the aqueous layer was extracted with ethyl acetate several times. The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered and evaporated to give the crude product, which was purified by flash chromatography (1% to 2% of ammonia saturated methanol in CH₂Cl₂) to give 2 (1.58 g; 52% yield).

B.



20


To a solution of 2 (1.54 g) in anhydrous THF (8 mL) was added a 1M solution of lithium aluminum hydride in diethyl ether (11 mL) at room temperature. The resulting solution was stirred for 2 hours and then diethyl ether (100 mL) was added. To the above mixture was carefully added a saturated aqueous sodium sulfate solution until no more hydrogen gas was evolved. The layers were separated, the aqueous layer was basified with K₂CO₃ to pH 9 and then extracted with ethyl acetate several times. The combined organic layers were washed

- 51 -

with brine, dried over anhydrous K_2CO_3 , and concentrated to give the crude product which was purified by flash chromatography (5% of ammonia saturated methanol in CH_2Cl_2) to give **3** (1.27 g; 85% yield).

5

C.




10

A solution of compound **3** (0.27 g) in a 0.5N HCl solution (20 mL) was heated to 90°C for 30 minutes. After the mixture was cooled to room temperature, the mixture was extracted with diethyl ether four times. The aqueous layer was separated and concentrated under vacuum to yield the crude product which was recrystallized from CH_3OH /diethyl ether to give **4** (0.141 g; 87% yield). MS (CI) 253 ($M + 1$).

15

EXAMPLE 2

A.




20

To a solution of 7 g of compound **1** in 35 mL of distilled water at 0°C was added 5 mL of a 50% aqueous NaOH solution. After 2 minutes of stirring, the mixture was extracted with methylene chloride. The combined organic extracts were washed with brine, dried over anhydrous Na_2SO_4 and concentrated. The residue was distilled under vacuum (40°C/18 mmHg) to give 2.7 g of free amine **2** (50% yield).

25


- 52 -

B.



To a mixture of magnesium turnings (0.73 g, 30.2 mmoles) in anhydrous THF (4 mL) at 50°C was added a solution of N-methyl-4-chloropiperidine **2** (2.7 g, 20.2 mmoles) in 2 mL of anhydrous THF over the course of 1.5 hours. After the addition was complete, the resulting mixture was heated to reflux for 1.5 hours and then it was cooled to -55°C. Copper (I) iodide (0.192 g) was then added and the resulting mixture was stirred at -55°C for 3.5 hours. A solution of N-trityl-4-chloromethylimidazole (4.83 g) in 20 mL of THF was added. The resulting mixture was stirred to -25°C for 1 hour, at room temperature for 45 minutes, and then it was cooled at -20°C. Aqueous saturated NH₄Cl solution was added to quench the reaction. The mixture was stirred at room temperature until the aqueous layer turned a deep blue color. The organic layer was separated and the aqueous layer was extracted with ethyl acetate. The combined organic solutions were washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated. The residue was purified by flash chromatography (MeOH/EtOAc) to yield 2.26 g (40% yield) of compound **3**.

C.



A solution of compound **3** (1.55 g) in 40 mL of 0.5N aqueous hydrochloric acid was heated to reflux for 30 minutes. The mixture was cooled to room temperature and extracted with diethyl ether. The aqueous solution was concentrated and the residue was


- 53 -

recrystallized from isopropanol/diethyl ether to give 0.6 g (65% yield) of compound **4**. MS (CI) 180 (M+1).


EXAMPLE 3

5

A.



B.




10

In a manner similar to that described in Example 2, Steps A and B, N-benzyl-4-chloropiperidine (**1**) is converted to compound **3**.

15

C.




20

To a solution of **3** (1.49 g) in 20 mL of anhydrous methanol was added 0.95g of ammonium formate (NH₄COO), and 1.49 g of 10% Pd/C. The mixture was refluxed under nitrogen for 40 minutes, cooled to room temperature, and then filtered through a pad of celite. The filtrate was concentrated, and the residue was purified by flash chromatography on silica gel to give 0.81 g (66% yield) of compound **4**.

- 54 -


D.



A solution of compound 4 (0.31 g) in 25 mL of 1N aqueous hydrochloric acid was heated to reflux for one hour. The mixture was cooled to room temperature and extracted with diethyl ether. The aqueous solution was concentrated and the residue was recrystallized from isopropanol/diethyl ether to give 0.16 g (64% yield) of compound 5. MS (CI) 166 (M + 1).


EXAMPLE 4

A.



15


To a mixture of magnesium turnings (0.73 g) in anhydrous THF (4mL) at 50°C was added a solution of N-methyl-4-chloropiperidine (2) (2.7 g) in 2 mL of anhydrous THF over the course of 1.5 hours. After the addition was complete, the resulting mixture was heated to reflux for an additional 1.5 hours and then the mixture was cooled to 0°C. A solution of N-tritylimidazoylcarboxaldehyde (1)



- 55 -

(6.83 g of 1 in 50 mL of THF) was added to the above Grignard reagent solution. The reaction mixture was stirred for 1 hour at 0°C and then quenched with water. The organic layer was separated and the aqueous layer was extracted with ethyl acetate. The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated. The residue was purified by flash chromatography on silica gel (eluting solvents: 1 to 10% ammonia saturated methanol in methylene chloride) to give 6.9 g (77% of compound 3).

10




A solution of 1.6 g of compound 3 in 40 mL of 85% sulfuric acid was stirred at room temperature for 24 hours. After the solution was cooled to 0°C, the reaction mixture was basified with KOH to pH = 9. The mixture was filtered and the filtrate was concentrated to approximately 1/3 volume. The resulting solution was extracted with 2:1 ethyl acetate/dichloromethane. The combined organic extracts were dried over anhydrous potassium carbonate, filtered, and concentrated to give an oil which was purified by preparative TLC (10% to 20% of ammonia saturated methanol in dichloromethane) to yield the free amine of 4. The free amine of 4 was dissolved in dilute HCl and pumped dry to give 0.303 g (33%) of compound 4. MS (CI) 178 (M+1).

25

The following are examples of pharmaceutical dosage forms which contain a compound of the invention. As used therein, the term "active compound" is used to designate the compound

- 56 -



- The scope of the invention in its pharmaceutical composition aspect is not to be limited by the examples provided, since
- 5 any other compound of structural formula 1.0 can be substituted into the pharmaceutical composition examples.

Pharmaceutical Dosage Form Examples

EXAMPLE A

Tablets

<u>No.</u>	<u>Ingredients</u>	<u>mg/tablet</u>	<u>mg/tablet</u>
1.	Active compound	100	500
2.	Lactose USP	122	113
3.	Corn Starch, Food Grade, as a 10% paste in Purified Water	30	40
4.	Corn Starch, Food Grade	45	40
5.	Magnesium Stearate	<u>3</u>	<u>7</u>
	Total	300	700

10

Method of Manufacture

- Mix Item Nos. 1 and 2 in a suitable mixer for 10–15 minutes. Granulate the mixture with Item No. 3. Mill the damp granules through a coarse screen (e.g., 1/4", 0.63 cm) if necessary. Dry the damp
- 15 granules. Screen the dried granules if necessary and mix with Item No. 4 and mix for 10–15 minutes. Add Item No. 5 and mix for 1–3 minutes.

SUBSTITUTE SHEET

- 57 -

Compress the mixture to appropriate size and weigh on a suitable tablet machine.

EXAMPLE B**Capsules**

5

<u>No.</u>	<u>Ingredient</u>	<u>mg/capsule</u>	<u>mg/capsule</u>
1.	Active compound	100	500
2.	Lactose USP	106	123
3.	Corn Starch, Food Grade	40	70
4.	Magnesium Stearate NF	<u>4</u>	<u>7</u>
	Total	250	700

Method of Manufacture


10 Mix Item Nos. 1, 2 and 3 in a suitable blender for 10-15 minutes. Add Item No. 4 and mix for 1-3 minutes. Fill the mixture into suitable two-piece hard gelatin capsules on a suitable encapsulating machine.

15 While the present invention has been described in conjunction with the specific embodiments set forth above, many alternatives, modifications and variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.

- 58 -

WHAT IS CLAIMED IS:

1. A compound of the formula:



or a pharmaceutically acceptable salt or solvate thereof, wherein:


- (A) m is an integer selected from the group consisting of: 1 and 2;
- (B) n and p are integers and are each independently selected from the group consisting of: 0, 1, 2, 3, and 4 such that the sum of n and p is 4 and T is a 6-membered ring;
- (C) R^3 and R^4 are each independently bound to the same or different carbon atom of ring T such that there is only one R^3 group and one R^4 group in ring T , and each R^1 , R^2 , R^3 , and R^4 is independently selected from the group consisting of:
- (1) H;
 - (2) C_1 to C_6 alkyl; and
 - (3) $-(CH_2)_q-R^6$ wherein q is an integer of: 1 to 7, and R^6 is selected from the group consisting of: phenyl, substituted phenyl, $-OR^7$, $-C(O)OR^7$, $-C(O)R^7$, $-OC(O)R^7$, $-C(O)NR^7R^8$, CN and $-SR^7$ wherein R^7 and R^8 are as defined below, and wherein the substituents on said substituted phenyl are each independently selected from the group consisting of: $-OH$, $-O-(C_1 \text{ to } C_6)\text{alkyl}$, halogen, C_1 to C_6 alkyl, $-CF_3$, $-CN$, and $-NO_2$, and wherein said substituted phenyl contains from 1 to 3 substituents;
- (D) R^5 is selected from the group consisting of:

- 59 -


- 5
- (1) H;
- (2) C₁ to C₂₀ alkyl;
- (3) C₃ to C₆ cycloalkyl;
- (4) -C(O)OR^{7'}; wherein R^{7'} is the same as R⁷ defined below except that R^{7'} is not H;
- (5) -C(O)R⁷;
- (6) -C(O)NR⁷R⁸;
- (7) allyl;
- (8) propargyl; and
- 10 (9) -(CH₂)_q-R⁶, wherein q and R⁶ are as defined above, and when q is equal to 1, then R⁶ is not OH or SH;
- (E) R⁷ and R⁸ are each independently selected from the group consisting of: H, C₁ to C₆ alkyl, and C₃ to C₆ cycloalkyl;
- (F) the dotted line (-----) represents a double bond that is optionally present when m is 1, and n is not 0, and p is not 0, and when said double bond is present then R² is absent; and
- 15 (G) when m is 2, each R¹ is the same or different substituent for each m, and each R² is the same or different substituent for each m, and at least two of the substituents R¹ and/or R² are H.
- 20

2. The compound of Claim 1 wherein said compound is selected from the group consisting of compounds having the formula:

25




- 60 -









5

wherein R^1 to R^5 are as defined for Formula 1.0.

3. The compound of Claim 1 wherein said compound
 10 is selected from the group consisting of compounds having the formula:



- 61 -



5

; and

wherein R¹ to R⁵ are as defined for Formula 1.0.

4. The compound of Claim 1 wherein R¹ to R⁴ are
 10 each independently selected from the group consisting of H, and C₁ to C₆ alkyl.

5. The compound of Claim 4 wherein m is 1.


15 6. The compound of Claim 4 wherein R⁵ is selected from the group consisting of: H, C₁ to C₆ alkyl, -C(O)OR⁷, -C(O)R⁷, -C(O)NR⁷R⁸, and -(CH₂)_q-R⁶ wherein R⁶ is phenyl.

- 62 -

7. The compound of Claim 6 wherein m is 1 and R² is H.

8. The compound of Claim 7 wherein R⁵ is selected from the group consisting of H and C₁ to C₆ alkyl, and R⁴ is H.

9. The compound of Claim 1 wherein said compound is selected from the group consisting of compounds having the formula:



wherein R¹ is selected from the group consisting of: H and C₁ to C₆ alkyl, and R⁵ is selected from the group consisting of: H, C₁ to C₆ alkyl, -C(O)OR⁶, -C(O)R⁷, -C(O)NR⁷R⁸, and -(CH₂)_q-R⁶ wherein R⁶ is phenyl.







- 63 -

10. The compound of Claim 9 wherein R⁵ is selected from the group consisting of: H and C₁ to C₆ alkyl.



11. The compound of Claim 10 wherein R¹ and R⁵ are independently selected from the group consisting of: H and methyl.

12. The compound of Claim 1 wherein said compound is selected from the group consisting of:


10



15



- 64 -




13. A pharmaceutical composition, for use in treating allergy, inflammation, hypertension, glaucoma, sleeping disorders, states of hyper and hypo motility of the gastrointestinal tract, hypo and hyperactivity of the central nervous system, Alzheimers, Schizophrenia, and migraines, comprising a pharmaceutically acceptable carrier and an effective amount of a Compound of Claim 1.
14. A method of treating allergy, inflammation, hypertension, glaucoma, sleeping disorders, states of hyper and hypo motility of the gastrointestinal tract, hypo and hyperactivity of the central nervous system, Alzheimers, Schizophrenia, and migraines comprising administering an effective amount of a compound of Claim 1 to a patient in need of such treatment.
15. The use of a compound of Claim 1 for the manufacture of a medicament for use in treating allergy, inflammation, hypertension, glaucoma, sleeping disorders, states of hyper and hypo motility of the gastrointestinal tract, hypo and hyperactivity of the central nervous system, Alzheimers, Schizophrenia, and migraines.
16. The use of a compound of Claim 1 for the treatment of allergy, inflammation, hypertension, glaucoma, sleeping disorders, states of hyper and hypo motility of the gastrointestinal tract, hypo and hyperactivity of the central nervous system, Alzheimers, Schizophrenia, and migraines.
17. A method of preparing a pharmaceutical composition comprising admixing a compound of Claim 1 with a pharmaceutically acceptable carrier.

- 65 -

18. A process for preparing a compound of Claim 1 comprising:

(A)




5

1) deprotecting compound (7) by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (8);

10

(B)



15

1) deprotecting compound (17), wherein R⁹ is an alkyl, cycloalkyl, benzyl, substituted benzyl, allyl or propargyl R⁵ group, following reaction path 6, by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (19); or

2) (a) treating compound (17), when R⁹ is not an R⁵


- 66 -

group, with tetrabutylammonium fluoride in tetrahydrofuran at a temperature of about 0 to about 50°C to produce compound (18);

(b1) reacting compound (18) with (i) R^5-X , when R^5 is $-C(O)R^7$, $-C(O)OR^{7'}$, $-C(O)NR^7R^8$ or alkyl, in an organic solvent optionally in the presence of a suitable base; or (ii) with $R^{5A}-CHO$, when R^5 is alkyl, cycloalkyl, allyl, propargyl, benzyl or substituted benzyl, under hydrogenating conditions, or in the presence of $NaBH_3(CN)$, in an organic solvent; wherein R^{5A} represents an R^5 group that has one less $-CH_2-$ group and X represents a suitable leaving group; said reaction (i) or (ii) being performed at a temperature within the range of about -30 to about 80°C; or

(b2) reacting compound (18) with $O=C=N-R^7$ in an inert organic solvent at a temperature in the range of about 20°C to reflux; and

(c) deprotecting the reaction product of (b1) or (b2) by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (19);




(a1) reacting compound (25) with (i) R^5-X , when R^5 is $-C(O)R^7$, $-C(O)OR^{7'}$, $-C(O)NR^7R^8$ or alkyl, in an organic solvent optionally in the presence of a suitable base; or (ii) with $R^{5A}-CHO$, when R^5 is alkyl, cycloalkyl, allyl, propargyl, benzyl or substituted benzyl, under hydrogenating conditions, or in the presence of $NaBH_3(CN)$, in an organic solvent; wherein R^{5A} represents an R^5 group that has one less

- 67 -

-CH₂- group and X represents a suitable leaving group; said reaction (i) or (ii) being performed at a temperature within the range of about -30 to about 80°C; or


(a2) reacting compound (25) with O=C=N-R⁷ in an inert organic solvent at a temperature in the range of about 20°C to reflux; and

(b) deprotecting the reaction product of (a1) or (a2) by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (26); or



deprotecting compound (24), wherein R is -CH₃ or -CH₂-C₆H₅, by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (26);

15




20 1) deprotecting compound (30), wherein R⁹ is an alkyl, cycloalkyl, benzyl, substituted benzyl, allyl or propargyl R⁵ group, by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (31); or

25 2) (a) treating compound (30), wherein R⁹ is a protecting group, with tetrabutylammonium fluoride in tetrahydrofuran at a temperature of about 0 to about 50°C;

- 68 -

- (b1) reacting the reaction product of (a) with (i) R^5-X , when R^5 is $-C(O)R^7$, $-C(O)OR^7$, $-C(O)NR^7R^8$ or alkyl, in an organic solvent optionally in the presence of a suitable base; or (ii) with $R^{5A}-CHO$, when R^5 is alkyl, cycloalkyl, allyl, propargyl, benzyl or substituted benzyl, under hydrogenating conditions, or in the presence of $NaBH_3(CN)$, in an organic solvent; wherein R^{5A} represents an R^5 group that has one less $-CH_2-$ group and X represents a suitable leaving group; said reaction (i) or (ii) being performed at a temperature within the range of about -30 to about $80^\circ C$; or
- (b2) reacting the reaction product of (a) with $O=C=N-R^7$ in an inert organic solvent at a temperature in the range of about $20^\circ C$ to reflux; and
- (c) deprotecting the reaction product of (b1) or (b2) by treatment with dilute aqueous acid at a temperature of about 70 to about $90^\circ C$ to produce compound (31);




- 1) deprotecting compound (36), wherein R^9 is an alkyl, cycloalkyl, benzyl, substituted benzyl, allyl or propargyl R^5 group, by treatment with dilute aqueous acid at a temperature of about 70 to about $90^\circ C$ to produce compound (26A); or
- 2) (a) treating compound (36), wherein R^9 is a protecting group, with tetrabutylammonium fluoride in tetrahydrofuran at a temperature of about 0 to about $50^\circ C$;
- (b1) reacting the reaction product of (a) with (i) R^5-X , when R^5 is $-C(O)R^7$, $-C(O)OR^7$, $-C(O)NR^7R^8$ or alkyl, in an organic solvent optionally in the presence of a suitable base; or (ii) with

- 69 -

- $R^{5A}-CHO$, when R^5 is alkyl, cycloalkyl, allyl, propargyl, benzyl or substituted benzyl, under hydrogenating conditions, or in the presence of $NaBH_3(CN)$, in an organic solvent; wherein R^{5A} represents an R^5 group that has one less $-CH_2-$ group and X represents a suitable leaving group; said reaction (i) or (ii) being performed at a temperature within the range of about -30 to about $80^\circ C$; or
- (b2) reacting the reaction product of (a) with $O=C=N-R^7$ in an inert organic solvent at a temperature in the range of about $20^\circ C$ to reflux; and
- (c) deprotecting the reaction product of (b1) or (b2) by treatment with dilute aqueous acid at a temperature of about 70 to about $90^\circ C$ to produce compound (26A);

(F)



15

- 1) (a) reacting compound (41) with (i) R^5-X , when R^5 is $-C(O)R^7$, $-C(O)OR^7$, $-C(O)NR^7R^8$ or alkyl, in an organic solvent optionally in the presence of a suitable base; or (ii) with $R^{5A}-CHO$, when R^5 is alkyl, cycloalkyl, allyl, propargyl, benzyl or substituted benzyl, under hydrogenating conditions, or in the presence of $NaBH_3(CN)$, in an organic solvent; wherein R^{5A} represents an R^5 group that has one less $-CH_2-$ group and X represents a suitable leaving group; said reaction (i) or (ii) being performed at a temperature within the range of about -30 to about $80^\circ C$ to produce compound (26A); or


20

25

- 70 -

(b) reacting compound (41) with $O=C=N-R^7$ in an inert organic solvent at a temperature in the range of about 20°C to reflux to produce compound (26A);


5



1) (a) reacting compound (19A') with (i) R^5-X , when R^5 is $-C(O)R^7$, $-C(O)OR^7$, $-C(O)NR^7R^8$ or alkyl, in an organic solvent optionally in the presence of a suitable base; or (ii) with $R^{5A}-CHO$, when R^5 is alkyl, cycloalkyl, allyl, propargyl, benzyl or substituted benzyl, under hydrogenating conditions, or in the presence of $NaBH_3(CN)$, in an organic solvent; wherein R^{5A} represents an R^5 group that has one less $-CH_2-$ group and X represents a suitable leaving group; said reaction (i) or (ii) being performed at a temperature within the range of about -30 to about 80°C to produce compound (19A); or

(b) reacting compound (19A') with $O=C=N-R^7$ in an inert organic solvent at a temperature in the range of about 20°C to reflux to produce compound (19A);

20




1) (a) reacting compound (8A') with (i) R^5-X , when R^5 is $-C(O)R^7$, $-C(O)OR^7$, $-C(O)NR^7R^8$ or alkyl, in an organic solvent optionally in the presence of a suitable base; or (ii) with $R^{5A}-CHO$, when

25

- 71 -

- R^5 is alkyl, cycloalkyl, allyl, propargyl, benzyl or substituted benzyl, under hydrogenating conditions, or in the presence of $\text{NaBH}_3(\text{CN})$, in an organic solvent; wherein R^{5A} represents an R^5 group that has one less $-\text{CH}_2-$ group and X represents a suitable leaving group; said reaction (i) or (ii) being performed at a temperature within the range of about -30 to about 80°C to produce compound (8A); or
- (b) reacting compound (8A') with $\text{O}=\text{C}=\text{N}-R^7$ in an inert organic solvent at a temperature in the range of about 20°C to reflux to produce compound (8A);


10



- 1) deprotecting compound (44') by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (44);

15

- 72 -




- 1) deprotecting compound (45''), wherein R⁹ is an alkyl, cycloalkyl, benzyl, substituted benzyl, allyl or propargyl R⁵ group, by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (45); or
- 2) (a) treating compound (45''), wherein R⁹ is a protecting group, with tetrabutylammonium fluoride in tetrahydrofuran at a temperature of about 0 to about 50°C to produce compound (45');
 - (b1) reacting compound (45') with (i) R⁵-X, when R⁵ is -C(O)R⁷, -C(O)OR^{7'}, -C(O)NR⁷R⁸ or alkyl, in an organic solvent optionally in the presence of a suitable base; or (ii) with R^{5A}-CHO, when R⁵ is alkyl, cycloalkyl, allyl, propargyl, benzyl or substituted benzyl, under hydrogenating conditions, or in the presence of NaBH₃(CN), in an organic solvent; wherein R^{5A} represents an R⁵ group that has one less -CH₂- group and X represents a suitable leaving group; said reaction (i) or (ii) being performed at a temperature within the range of about -30 to about 80°C; or
 - (b2) reacting compound (45') with O=C=N-R⁷ in an inert organic solvent at a temperature in the range of about 20°C to reflux; and
 - (c) deprotecting the reaction product of (b1) or (b2)

- 73 -

by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (45);

(K)



5

- 1) deprotecting compound (47''), wherein R⁹ is an alkyl, cycloalkyl, benzyl, substituted benzyl, allyl or propargyl R⁵ group, by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (47); or
- 2) (a) treating compound (47''), wherein R⁹ is a protecting group, with tetrabutylammonium fluoride in tetrahydrofuran at a temperature of about 0 to about 50°C to produce compound (47');
 (b1) reacting compound (47') with (i) R⁵-X, when R⁵ is -C(O)R⁷, -C(O)OR⁷, -C(O)NR⁷R⁸ or alkyl, in an organic solvent optionally in the presence of a suitable base; or (ii) with R^{5A}-CHO, when R⁵ is alkyl, cycloalkyl, allyl, propargyl, benzyl or substituted benzyl, under hydrogenating conditions; or in the presence of NaBH₃(CN), in an organic solvent; wherein R^{5A} represents an R⁵ group that has one less -CH₂- group and X represents a suitable leaving group; said reaction (i) or (ii) being performed at a temperature within the range of about -30 to about 80°C; or


15

20

- 74 -

(b2) reacting compound (47') with $O=C=N-R^7$ in an inert organic solvent at a temperature in the range of about 20°C to reflux; and

- (c) deprotecting the reaction product of (b1) or (b2) by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (47);




- (a1) reacting compound (51') with (i) R^5-X , when R^5 is $-C(O)R^7$, $-C(O)OR^7$, $-C(O)NR^7R^8$ or alkyl, in an organic solvent optionally in the presence of a suitable base; or (ii) with $R^{5A}-CHO$, when R^5 is alkyl, cycloalkyl, allyl, propargyl, benzyl or substituted benzyl, under hydrogenating conditions, or in the presence of $NaBH_3(CN)$, in an organic solvent; wherein R^{5A} represents an R^5 group that has one less $-CH_2-$ group and X represents a suitable leaving group; said reaction (i) or (ii) being performed at a temperature within the range of about -30 to about 80°C; or


(a2) reacting compound (51') with $O=C=N-R^7$ in an inert organic solvent at a temperature in the range of about 20°C to reflux; and

- (b) deprotecting the reaction product of (a1) or (a2) by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (51); or

- 75 -



- deprotecting compound (51''), wherein R is -CH₃ or -CH₂-C₆H₅, by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (51);



10

- 1) deprotecting compound (52'), wherein R⁹ is an alkyl, cycloalkyl, benzyl, substituted benzyl, allyl or propargyl R⁵ group, by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (52); or

15

- 2) (a) treating compound (52'), wherein R⁹ is a protecting group, with tetrabutylammonium fluoride in tetrahydrofuran at a temperature of about 0 to about 50°C;

20

- (b1) reacting the reaction product of (a) with (i) R⁵-X, when R⁵ is -C(O)R⁷, -C(O)OR⁷, -C(O)NR⁷R⁸ or alkyl, in an organic solvent optionally in the presence of a suitable base; or (ii) with R^{5A}-CHO, when R⁵ is alkyl, cycloalkyl, allyl, propargyl, benzyl or substituted benzyl, under hydrogenating conditions, or in the presence of NaBH₃(CN), in an organic solvent; wherein R^{5A} represents an R⁵

- 76 -


group that has one less $-\text{CH}_2-$ group and X represents a suitable leaving group; said reaction (i) or (ii) being performed at a temperature within the range of about -30 to about 80°C ; or

(b2) reacting the reaction product of (a) with

- 5 $\text{O}=\text{C}=\text{N}-\text{R}^7$ in an inert organic solvent at a temperature in the range of about 20°C to reflux; and

(c) deprotecting the reaction product of (b1) or (b2) by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compound (52);

10




- 1) converting compound (53) to compound (54) using
- 15 an acidic aqueous solution and a temperature of about 50 to about 100°C , wherein $\text{R}^{5'}$ is an H, alkyl, cycloalkyl, benzyl, substituted benzyl, allyl or propargyl group; or
- 2) (a) reacting compound (54), wherein $\text{R}^{5'}$ is an H, with (i) R^5-X , when R^5 is $-\text{C}(\text{O})\text{R}^7$, $-\text{C}(\text{O})\text{OR}^7$, $-\text{C}(\text{O})\text{NR}^7\text{R}^8$ or alkyl, in an organic solvent optionally in the presence of a suitable base; or (ii) with $\text{R}^{5\text{A}}-\text{CHO}$, when R^5 is alkyl, cycloalkyl, allyl, propargyl, benzyl or substituted benzyl, under hydrogenating conditions, or in the presence of
- 20 $\text{NaBH}_3(\text{CN})$, in an organic solvent; wherein $\text{R}^{5\text{A}}$ represents an R^5 group

- 77 -

that has one less $-CH_2-$ group and X represents a suitable leaving group; said reaction (i) or (ii) being performed at a temperature within the range of about -30 to about 80°C , to produce compound (55); or


- (b) reacting compound (54), wherein R^5 is an H, with $O=C=N-R^7$ in an inert organic solvent at a temperature in the range of about 20°C to reflux, to produce compound (55);



10


- 1) deprotecting compounds (64') and (64A') by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compounds (64) and (64A), respectively;

- 78 -




- 1) deprotecting compounds (66') and (66A') by treatment with dilute aqueous acid at a temperature of about 70 to about 90°C to produce compounds (66) and (66A), respectively; or
- (Q) Preparing compounds of Claim 1, wherein R⁵ is H, by reacting an intermediate compound of one of the above process steps (A), (B), (C), (D), (E), (F), (G), (H), (I), (J), (K), (L), (M), (N), (O), or (P) with aqueous acid, at a temperature of about 25 to about 100°C, said intermediate compound having the imidazole nitrogen protected by Z and having the nitrogen of the cyclic six membered amine substituted with -C(O)O(t-butyl) or unsubstituted.

19. A compound of the formula:



20. A compound of the formula:



INTERNATIONAL SEARCH REPORT

PCT/US 92/10698

International Application No.

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C07D401/06; A61K31/415		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C07D ; A61K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	EP,A,0 197 840 (INSTITUT NATIONAL DE LA SANTE ET DE LA RECHERCHE MEDICALE (INSERM)) 15 October 1986 cited in the application page 18, first paragraph see claim 1	1-13, 15, 17, 18
A	US,A,4 925 851 (HOULIHAN) 15 May 1990 cited in the application formula I	1-13, 15, 17, 18
A	US,A,4 767 778 (ARRANG) 30 August 1988 cited in the application formula I	1-13, 15, 17, 18
A	EP,A,0 372 125 (AKZO) 13 June 1990 cited in the application formula I	1-13, 15, 17, 18
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
15 MARCH 1993	23. 03. 93	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	GETTINS M.P.	

Form PCT/ISA/210 (second sheet) (January 1985)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 92/ 10698

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 14, 16
because they relate to subject matter not required to be searched by this Authority, namely:
Please see Rule 39.1(iv) - PCT:

Methods for treatment of the human or animal body by surgery or therapy,
as well as diagnostic methods
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such
an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9210698
SA 68305

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 15/03/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0197840	15-10-86	FR-A- 2579596 JP-A- 61267574 US-A- 4707487	03-10-86 27-11-86 17-11-87
US-A-4925851	15-05-90	None	
US-A-4767778	30-08-88	FR-A- 2586562 DE-A- 3685170 EP-A,B 0214058 JP-A- 62123174	06-03-87 11-06-92 11-03-87 04-06-87
EP-A-0372125	13-06-90	AU-B- 623942 AU-A- 4440689 EP-A- 0368384 JP-A- 2180881 US-A- 5128340	28-05-92 17-05-90 16-05-90 13-07-90 07-07-92

EPO FORM P0479

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82